

IJMA



INTERNATIONAL JOURNAL OF MEDICAL ARTS

Volume 7, Issue 10 (October 2025)



<http://ijma.journals.ekb.eg/>

P-ISSN: 2636-4174

E-ISSN: 2682-3780



Available online at Journal Website
<https://ijma.journals.ekb.eg/>
 Main Subject [Orthopedic Surgery]



Original Article

Treatment of Supracondylar Humerus Fracture in Adults by Parallel Plates

Ahmed Mohamed Elhady Khater^{*1}; Samir Ahmed El-Shoura²; Mohamed Abdalah Hassan²

¹ Department of Orthopedic Surgery, Damietta General Hospital, Ministry of Health, Damietta, Egypt.

² Department of Orthopedic Surgery, Damietta Faculty of Medicine, Al-Azhar University, Damietta, Egypt.

Abstract

Article information

Received: 12-07-2023

Accepted: 18-04-2025

DOI: [10.21608/ijma.2024.222729.1740](https://doi.org/10.21608/ijma.2024.222729.1740).

*Corresponding author

Email: drhaady3@gmail.com

Citation: Khater AME, El-Shoura SA, Hassan MA. Treatment of Supracondylar Humerus Fracture in Adults by Parallel Plates. IJMA 2025 October; 7[10]: 6193-9198. doi: [10.21608/ijma.2024.222729.1740](https://doi.org/10.21608/ijma.2024.222729.1740).

Objective: The aim of this study was to evaluate the outcome of parallel plates for the treatment of supracondylar humerus fractures in adults.

Patients and Methods: This study included 20 adult patients with supracondylar humerus fractures. They were treated by open reduction and internal fixation by parallel plates through olecranon osteotomy technique. They were followed up clinically and radiologically for 6 months. On admission, all were assessed by careful history taking and detailed clinical examination. The preoperative radiological workup was performed by antero-posterior and lateral radiographs of the elbow with traction and computed tomography was performed with three-dimensional reconstruction. Postoperative, patients were assessed after 2, 4, 8, 12 weeks and 6 months of surgery, with clinical and radiological evaluations and assessed for pain, swelling, range of joint motion and radiological union. The functional assessment of the patient was done according to Mayo elbow performance index and the outcome was graded into excellent, good, poor and fair.

Results: The excellent outcome was achieved among 55.0%, while 30.0% had good, 10% had fair and 5.0% had poor outcome. The overall complication rate was 30.0% and the commonest were pain and stiffness [15.0% for each]. The mean±SD times to clinical and radiological union were 12.75 ± 6.44 and 14.70 ± 7.84 weeks, respectively. The excellent outcome was significantly associated with lower rate of complications and shorter time to clinical and radiological union.

Conclusion: Parallel plates for the treatment of adult supracondylar humerus fractures could be considered safe and effective technique, leading to good functional outcome and a complication rate comparable to published literature.

Keywords: Fracture fixation; Humeral Fractures; Elbow; Olecranon; Parallel Plates.



This is an open-access article registered under the Creative Commons, ShareAlike 4.0 International license [CC BY-SA 4.0] [<https://creativecommons.org/licenses/by-sa/4.0/legalcode>].

INTRODUCTION

Fracture distal humerus accounts for 2 -6 % of all the fractures and its incidence in the adults around 5.7 per 100,00 per year [1,2]. In young patients, distal humeral fractures are commonly caused by high velocity injuries, such as sports injuries or road traffic accidents. In contrast, distal humeral fractures in elderly people are predominantly low velocity injuries complicated by poor bone quality [3]. Distal humerus fracture can be classified based on the Orthopedic Trauma Association / *Arbeitsgemeinschaft für Osteosynthesefragen* [OTA/AO] classification system as extra-articular fractures [type A], partial articular fractures [type B], and complete articular fractures [type C]. Intra-articular fractures incorporate trochlea and capitulum fractures [4].

The examining surgeon must play out an exhaustive neurovascular assessment with all speculated elbow fractures; perceive subtle fracture; give sufficient immobilization and survey whether fractures require admission, quick orthopedic assessment or less urgent referral [5,6]. Dealing with such fractures can be very challenging due to the intricate anatomy of the elbow and small fracture fragments. In adults, the gold standard treatment for the distal humerus fractures is considered to be the open reduction internal fixation [ORIF] with plate fixation of both columns [7,8].

Open reduction and internal fixation with plates with good articular reconstruction have demonstrated satisfactory clinical outcomes. Firm stabilization can be achieved through various methods of fixation of plates [9-11]. Among them, several mechanical studies have proved that double plate fixation provides more stable fixation than other methods [12-14]. Various plate designs have been developed for the fixation of these fractures, some are like Y-plates, recon plates, precontoured anatomical plates. However, controversy still exists concerning the plate positions in terms of providing optimal stability for distal humerus fractures. The most widely used plate fixation method is placing plates perpendicular to each other, one over medial supracondylar ridge and another one over the flat posterior surface of lateral column called orthogonal plating. Stoffel et al demonstrated on mechanical studies two plates placed parallel to each other, one over each supracondylar ridge, providing better stability in compression and external rotation than to perpendicular plating system in cadaveric models [15-17].

AIM OF THE WORK

The aim of this study was to evaluate the outcome of parallel plates for the treatment of supracondylar humerus fractures in adults.

PATIENTS AND METHODS

This clinical trial had been conducted for supracondylar humerus fractures in adult patients admitted to the department of Orthopedic Surgery, Al-Azhar University Hospital [New Damietta]. Patients were treated by open reduction and internal fixation by parallel plates during the period from March to September 2022. They were followed up clinically and radiologically for 6 months.

The inclusion criteria: 1] Recent supracondylar humerus fractures, 2] Age between 18 and 60 years, and 3] Isolated closed supracondylar traumatic humerus fractures. On the other side, the exclusion criteria were the presence of open injuries, polytrauma patients, pathological fractures [except osteoporosis], and fractures with distal neurovascular injury.

Sampling technique: A non-probability [convenient] sample technique was used to recruit cases. All patients admitted in the Orthopedic Department during the study period and who's matching the inclusion criteria were recruited to participate in this study after taking the informed consent. They were [20] patients.

Ethical considerations: The details of the operation technique & complications were explained to the patients and informed written consent was obtained. The right to withdraw was granted and the collected data were obtained for research purposes only without divulging any data talk about the personality of the patient [Confidentiality will be ascertained]. In addition, the study protocol approved by the local research and Ethics Committee of Al-Azhar Faculty of Medicine [New Damietta] [IRB# DFM-IRB00012367-22-03-008].

All patients were treated with open reduction and internal fixation with parallel plates through olecranon osteotomy technique, and they were followed up clinically and radiologically for 6 months. On admission, all were assessed by careful history taking and detailed clinical examination. This included inquiry about personal data, comorbid disease conditions [e.g., diabetes mellitus [DM], Cardiac disease, hypertension, etc.], the mechanism of injury, pain or swelling, deformity, wounds, ecchymosis, neurovascular integrity, or associated injuries. The neurological examination was performed to assess the ulnar, radial and median nerve function. The preoperative radiological assessment was performed by antero-posterior and lateral radiographs of the elbow with traction. Computerized tomography of the elbow was performed especially in partial or very distal fractures because the various fragments usually superimposed, which hinders precise analysis of the fracture on standard views. Three-dimensional reconstruction showed the shape and position of the bone fragments and was helpful in determining the appropriate surgical approach.

To complete preoperative evaluation, laboratory tests were performed. This included complete blood count and bleeding profile, kidney and liver function tests, fasting blood sugar and postprandial blood sugar tests. The preoperative preparation achieved by the application of the above elbow slab with good padding to all the bony prominences, proper analgesia, control of any comorbid condition, and intravenous antibiotic administration, one hour before surgery. An informed consent for surgery was signed by the patient and his first close relative. This was signed after full explanation of the surgery and possible complications.

All surgeries were performed under general anesthesia in lateral decubitus position with arm supported and forearm hanging. The elbow was then exposed through standard midline posterior approach with incision beginning 5cm distal to the tip of the olecranon and extending proximally in the arm up to 10 cm above the tip of the olecranon [Figure 1]. The ulnar nerve was exposed and secured [Figure 2]. Ulnar nerve was identified proximally along the medial border of triceps, then released through the cubital tunnel down until the first motor branch by incision of the flexor pronator aponeurosis as the nerve passes between the two heads of flexor carpi ulnaris [FCU]. The site of the osteotomy was determined by subperiosteal reflection of the anconeus laterally and the ulna head of FCU medially to expose the olecranon. The bone was cleared with a small elevator at the site of the planned osteotomy. About 1.5 cm from the tip of olecranon the apex distal chevron osteotomy was performed with an oscillating saw and completed with an osteotome to facilitate interdigitation. A 'V' shaped Olecranon osteotomy was done to get better exposure of the articular surface in all cases as proximal mobilization of the osteotomized fragment and triceps allows ample

exposure of the articular surface and columns [Figure 3]. The articular fragments of intercondylar humerus were reduced and held with 'K' wires and then fixed with intercondylar screw. The reconstituted articular [condylar] block was reduced to a more stable column, and one or more K-wires were used for preliminary fixation. The comminuted fragments of the other column were reduced into correct alignment, and one or more K-wires were used for preliminary fixation. The accurate alignment of the articular block to the shaft was assured. One third tubular plates were placed parallel to each other on medial and lateral columns [Figure 4]. The plates were secured with 3.5 mm cortical screws in diaphyseal area and with 4 mm cancellous screws in metaphyseal area. Proximal end of both plates was not at the same level

to avoid stress riser fracture. The stability of the internal fixation was tested by moving the elbow through full range of motion. The Olecranon osteotomy was reduced and held with pointed reduction forceps to apply compression. Then distal to the fracture line, a hole was drilled through the ulna. Then a tension band was inserted through a drilled hole. then 2 k-wires were inserted through the proximal end of olecranon towards the anterior cortex under direct vision. The tension band was passed under triceps tendon in a figure of '8" around 2-pointed k-wires [Figure 5]. After fixation of the osteotomy the elbow was again put through the range of motion to test the stability of fixation. Suction drain was inserted, and incision was closed in layers by sutures or surgical staples. The arm was bandaged to support and protect the surgical wound.



Figure [1]: Incision over the elbow



Figure [2]: Ulnar nerve exploration



Figure [3]: V-shaped olecranon osteotomy




Figure [4]: K-wire and parallel plates fixation



Figure [5]: K- wire and tension band of olecranon




Figure [6]: X ray AP and Lateral of elbow after fixation

In the postoperative period, the limb was kept elevated. Active movement of fingers and elbow joint were encouraged from second

postoperative day. Suction drain was removed after 24 hours. Antibiotic prophylaxis was given as intravenous administration for 3 days and

continued with oral antibiotic for 4 days. Analgesics were given until the pain was subsided. Suture/staples were removed on the 14th postoperative day. Post-operative radiographs were obtained as soon as patient was comfortable [Figure 6]. Elbow was mobilized through full range of movement at least twice daily and patient was discharged with instruction to carry out physiotherapy outside the hospital in the form of active flexion-extension and pronation-supination exercises without loading. Patient was assessed after 2, 4, 8, 12 weeks and 6 months of surgery. At every follow up a detailed clinical examination was done and patient was assessed subjectively for the pain, swelling, range of joint motion and radiological union. The functional assessment of the patient was done according to Mayo elbow performance index.

The Mayo Elbow Performance score [MEPS] is an instrument used to test the limitations of the elbow during activities of daily living [ADL] [18]. It included 4 domains [pain, range of motion, stability and daily function]. The total score reached 100 and value below 60 was considered poor. Values between 60 and 74 are fair, while values between 75 and 89 are good, and values between 90 and 100 are excellent.

Statistical analysis: Data was coded and entered to a Microsoft Excel Sheet. Data were then imported into Statistical Package for Social Sciences [SPSS] version 16.0 [SPSS Inc., Chicago, USA] for analysis. The relative frequencies and percentages were calculated for qualitative

[categorical variables], while mean, standard deviation, minimum and maximum values were computed for quantitative continues variables.

RESULTS

The current study included 20 patients, their age ranged between 28 and 50 years, with male-sex predominance [males represented 70.0%]; smoking practiced by 35%, while 20.0% had hypertension and 15% had diabetes mellitus. The road traffic accident was the major mode of injury [75.0%], with right predominance [70.0%] and 90% of subjects were right-handed [Table 1].

According to MEPS grading system, 55.0% had an excellent outcome, 30.0% had good, 10% had fair and 5.0% had poor outcome. The complications were reported among 30.0% of subjects and commonest were pain and stiffness [15.0% for each complication]. The mean±SD times to clinical and radiological union were 12.75 ± 6.44 and 14.70 ± 7.84 weeks, respectively [Table 2]. The excellent outcome when compared to other outcomes was significantly associated with significantly lower rate of complications and stiffness. In addition, it is significantly associated with shorter time to clinical and radiological union. Otherwise, no significant association was discovered [Table 3]. The laboratory data was mainly in normal values and showed no significant association with excellent outcome [Data not tabulated].

Table [1]: Demographic data of the studied patients.

Variables		Statistics
Age [year]	Mean ±SD	37.45±6.23
	Min. – Max.	28- 50
Sex [n, %]	Male	14 [70.0%]
	Female	6 [30.0%]
Comorbid conditions [n, %]	Smoking	7 [35.0%]
	Hypertension	4 [20.0%]
	Diabetes mellitus	3 [15.0%]
Mode of injury [n, %]	RTA	15 [75.0%]
	Fall from height/heavy object	3 [15.0%]
	Assault	2 [10.0%]
Injured side [n, %]	Right	14 [70.0%]
	Left	6 [30.0%]
Dominant hand [n,%]	Right	18 [90.0%]
	Left	2 [10.0%]

Table [2]: Outcome among the study subjects.

		Statistics
MEPS Outcome	Excellent	11 [55.0%]
	Good	6 [30.0%]
	Fair	2 [10.0%]
	Poor	1 [5.0%]
Postoperative complications [n, %]	Total complications	6 [30.0%]
	Myositis	1 [5.0%]
	Stiffness	3 [15.0%]
	Infection	2 [10.0%]
	Non-union at osteotomy	1 [5.0%]
	Pain	3 [15.0%]
	Hardware prominence	1 [5.0%]
Time to	Clinical union [weeks]	12.75 ± 6.44
	Radiological union [weeks]	14.70 ± 7.84
Mayo score		86.35 ± 12.67

Table [3]: Factors associated with excellent outcome compared to others

		Excellent outcome [n=11]	Other outcome grades [n=9]	Test	p
Age [years]		37.90±6.65	36.89±6.03	0.35	0.72
Sex	Male	9 [81.8%]	5 [55.6%]	1.62	0.20
	Female	2 [18.2%]	4 [44.4%]		
Comorbid conditions	Smoking	4 [36.4%]	3 [33.3%]	0.02	0.89
	Hypertension	2 [18.2%]	1 [11.1%]	0.19	0.66
	Diabetes mellitus	3 [27.3%]	1 [11.1%]	0.81	0.36
Mode of injury	RTA	8 [72.7%]	7 [77.8%]	2.22	0.32
	Fall from height/heavy object	1 [9.1%]	2 [22.2%]		
	Assault	2 [18.2%]	0 [0.0%]		
Injured side	Right	8 [72.7%]	6 [66.7%]	0.09	0.76
	Left	3 [27.3%]	3 [33.3%]		
PO complications	Total complications	1 [9.1%]	5 [55.6%]	5.08	0.024*
	Myositis	0 [0.0%]	1 [11.1%]	1.28	0.25
	Stiffness	0 [0.0%]	3 [33.3%]	4.31	0.038*
	Infection	0 [0.0%]	2 [22.2%]	2.71	0.10
	Non-union at osteotomy	0 [0.0%]	1 [11.1%]	1.28	0.25
	Pain	1 [9.1%]	2 [22.2%]	0.67	0.41
	Hardware prominence	0 [0.0%]	1 [11.1%]	1.28	0.25
Time to	Clinical union [weeks]	9.18±1.60	17.11±7.52	3.42	0.003*
	Radiological union [weeks]	10.91±1.36	19.33±9.35	2.97	0.008*

DISCUSSION

This study was conducted for supracondylar humerus fracture in adults treated by open reduction and internal fixation by parallel plates by olecranon osteotomy technique and they were followed up clinically and radiologically for 6 months.

Yadav et al. [19] reported that, the olecranon osteotomy approach does not seem to have a clinical advantage over the approaches that preserve the integrity of the elbow extensor device. However, a systematic review and meta-analysis conducted by **Chen et al.** [20] compared various surgical approaches on elbow functional outcomes for patients with distal humerus intercondylar fractures [DHF]. They concluded olecranon osteotomy was superior than triceps-sparing approach in restoring joint function. The meta-analysis included four studies with 276 patients and compared olecranon osteotomy with triceps-sparing. The pooled analysis indicated that patients treated using olecranon osteotomy had a better functional outcome than patients treated with triceps-sparing. Thus, we used it on the current work.

Results of the current work are in line with **Kumar et al.** [21] who included 23 subjects, their mean age was 39.1±11.5 years with male sex predominance [69.6%]. They reported union rate in all patients and complications rate of less than 19%. However, they followed up their patients for a longer duration [for up to one year] that the current study.

Salvador et al. [22] treated 27 supracondylar humeral fractures and achieved 96.0% complete union at 6 months of follow up. They reported no infection. However, two subjects needed another surgery to treat stiff elbow. 60% of subjects were able to return to their previous usual activities. To explain their results in the light of the current one, they included more older patients [the mean age was 56 ± 22.9 years] with higher females [52.0%].

In addition, **Shah et al.** [23] reported excellent outcome in 41 [56.94%], good in 21 [29.17%], fair in 6 [8.33%] and poor outcome in 4 [5.56%] of their patients. These results are comparable to the current study. However, **Singh et al.** [24] reported that total of 25 patients, 8 patients [32%] got excellent results, 14 patients [56%] got good outcome and 3 [12%] got poor outcomes.

Most recently, **Jagadish U et al.** [25] reported on a total of 30 patients with distal humerus fracture fixed with parallel plates. The excellent outcome was reported among 47%, while 33.0% had good outcome, 13.0% had fair and 7.0% had poor outcome. The results which comparable to the current work.

Postoperative pain was reported among 15.0% in the current work. However, **Salvador et al.** [22] reported mild and moderate pain in 9 and 6 out of 27 patients included in their study, which is quite different and higher than the current work. This could be explained by the older age of their patients and the predominance of females, where the threshold of pain is lower among them than younger [adult] males [the predominant in the current work].

Additionally, **Patel et al.** [26] reported a complication rate of 61%. Among all patients, 49% required a reoperation. Elbow stiffness [19%] was the commonest complication followed by nerve palsy [16%]. The non-union rate was 9.0% and the deep infection, painful implants, post-traumatic arthritis and heterotopic ossification [9% for each of them]. The higher rate and different distribution of complications are due to heterogenous surgical procedures according to surgeon preferences. Furthermore, **Singh et al.** [24] revealed a complication rate of 44%. The commonest were painful hardware, superficial infection and transient ulnar nerve palsy.

Conclusion: Parallel plates for the treatment of adult supracondylar humerus fractures could be considered safe and effective technique, leading to good functional outcome and lower complication rate. However, the relatively low number of patients, shorter duration of follow up, absence of comparative groups or procedures are limiting steps of the current work. This advocate future studies including higher number of patients with longer duration of follow up. The current one represented one step on the road to determine the ideal intervention.

Financial and non-financial activities and relationships of interest: None

REFERENCES

1. Beazley JC, Baraza N, Jordan R, Modi CS. Distal Humeral Fractures-Current Concepts. *Open Orthop J*. 2017 Nov 30; 11:1353-1363. doi: 10.2174/1874325001711011353.
2. Vauclair F, Goetti P, Nguyen NTV, Sanchez-Sotelo J. Distal humerus nonunion: evaluation and management. *EFORT Open Rev*. 2020 May 1;5[5]:289-298. doi: 10.1302/2058-5241.5.190050.
3. Lan X, Zhang LH, Tao S, Zhang Q, Liang XD, Yuan BT, Xu WP, Yin P, Tang PF. Comparative study of perpendicular versus parallel double plating methods for type C distal humeral fractures. *Chin Med J [Engl]*. 2013 Jun;126[12]:2337-42. PMID: 23786950.
4. Meinberg E, Agel J, Kellam JF, Roberts CS. The Fracture and Dislocation Classification Compendium 2017 - "Nearing the Finish Line". *Injury*. 2017 Apr;48[4]:793-794. doi: 10.1016/j.injury.2017.03.032.
5. Goldflam K. Evaluation and treatment of the elbow and forearm injuries in the emergency department. *Emerg Med Clin North Am*. 2015 May;33[2]:409-21. doi: 10.1016/j.emc.2014.12.010.
6. Hanlon DP, Mavrophilipos V. The Emergent Evaluation and Treatment of Elbow and Forearm Injuries. *Emerg Med Clin North Am*. 2020 Feb;38[1]:81-102. doi: 10.1016/j.emc.2019.09.005.
7. Tyllianakis M, Solou K, Lakoumentas J, Panagopoulos A. Long-Term Functional Outcomes and Complications of Intra-Articular [AO type B, C] Distal Humerus Fractures in Adults: A Retrospective Review. *Cureus*. 2022 Jan 10;14[1]: e21094. doi: 10.7759/cureus.21094.
8. Spierings KE, Schoolmeesters BJ, Doornberg JN, Eygendaal D, van den Bekerom MP. Complications of olecranon osteotomy in the treatment of distal humerus fracture. *Clin Shoulder Elb*. 2022 Jun;25[2]:163-169. doi: 10.5397/cise. 2021.00591.
9. Jacquot A, Poussange N, Charrissoux JL, Clavert P, Obert L, Pidhorz L, Sirveaux F, Mansat P, Fabre T. Usefulness and reliability of two- and three-dimensional computed tomography in patients older than 65 years with distal humerus fractures. *Orthop Traumatol Surg Res*. 2014 May;100[3]:275-80. doi: 10.1016/j.otsr.2014.01.003.
10. Cañete San Pastor P, Lopez Valenciano J, Copete I, Prosper Ramos I. Fixation of olecranon osteotomy only with 6.5 mm partially threaded cancellous screw is a safe and effective method used in surgical management of distal humerus fractures. *J Exp Orthop*. 2021 Jan 12;8[1]:4. doi: 10.1186/s40634-020-00317-8.
11. Sanchez-Sotelo J, Torchia ME, O'Driscoll SW. Complex distal humeral fractures: internal fixation with a principle-based parallel-plate technique. Surgical technique. *J Bone Joint Surg Am*. 2008 Mar;90 Suppl 2 Pt 1:31-46. doi: 10.2106/JBJS.G.01502.
12. Got C, Shuck J, Biercevicz A, Paller D, Mulcahey M, Zimmermann M, Blaine T, Green A. Biomechanical comparison of parallel versus 90-90 plating of bicondylar distal humerus fractures with intra-articular comminution. *J Hand Surg Am*. 2012 Dec;37[12]:2512-8. doi: 10.1016/j.jhsa.2012.08.042.
13. Wang X, Liu G. A comparison between perpendicular and parallel plating methods for distal humerus fractures: A meta-analysis of randomized controlled trials. *Medicine [Baltimore]*. 2020 Jun 5;99[23]:e19602. doi: 10.1097/MD.00000000000019602.
14. Zalavras CG, Papasoulis E. Intra-articular fractures of the distal humerus-a review of the current practice. *Int Orthop*. 2018 Nov;42[11]:2653-2662. doi: 10.1007/s00264-017-3719-4.
15. Athanasiou ED, Komnos G, Deligeorgis D, Hantes M, Karachalios T, Malizos KN, Varitimidis S. Double Plating in Type C Distal Humerus Fractures: Current Treatment Options and Factors that Affect the Outcome. *Strategies Trauma Limb Reconstr*. 2022 Jan-Apr;17[1]:7-13. doi: 10.5005/jp-journals-10080-1546.
16. Haglin JM, Kugelman DN, Lott A, Belayneh R, Konda SR, Egol KA. Intra-articular Distal Humerus Fractures: Parallel Versus Orthogonal Plating. *HSS J*. 2022 May;18[2]:256-263. doi: 10.1177/15563316211009810.
17. Chen C, Xiao D, Li T, Gong M, Zha Y, Hua K, Sun W, Ji S, Jiang X. Comparative study of functional outcomes between OTA/AO type C, Gustilo type I/II open fractures and closed fractures of the distal humerus treated by open reduction and internal fixation. *BMC Musculoskelet Disord*. 2021 Nov 10;22[1]:939. doi: 10.1186/s12891-021-04817-1.
18. Schneeberger AG, Kösters MC, Steens W. Comparison of the subjective elbow value and the Mayo elbow performance score. *J Shoulder Elbow Surg*. 2014;23[3]:308-12. doi: 10.1016/j.jse.2013.11.018.
19. Yadav V, Sharma P, Gohiya A. Functional outcome of intraarticular distal humerus fracture fixation using triceps-sparing paratricipital approach. *Indian J Orthop*. 2016;50[6]:595-601. doi: 10.4103/0019-5413.193487.
20. Chen H, Li D, Zhang J, Xiong X. Comparison of treatments in patients with distal humerus intercondylar fracture: a systematic review and meta-analysis. *Ann Med*. 2017;49[7]:613-625. doi: 10.1080/07853890.2017.1335429.
21. Kumar S, Singh S, Kumar D, Kumar N, Verma R. Intercondylar humerus fracture- parallel plating and its results. *J Clin Diagn Res*. 2015 Jan;9[1]:RC01-4. doi: 10.7860/JCDR/2014/12137.5479.
22. Salvador J, Castellón P, Fuentes I, Bernaus M, Anglès F. Double plating as treatment for supracondylar humeral fractures. *Rev Esp Cir Ortop Traumatol*. 2017 Sep-Oct;61[5]:324-330. doi: 10.1016/j.recot.2017.05.003.
23. Shah SW, Bashir T, Aslam MU. Functional Outcomes of Pre-Contoured Parallel Plates in Distal Humerus AO/OTA Type C Fractures in adults. *PJMHS* 2020; 14 [4]: 906-908.
24. Singh D, Sareen AS, Sandhu A. Comparative study of functional outcome of distal humerus intra-articular fracture treated with parallel plates and with orthogonal plates [A study of 30 cases]. *European Journal of Molecular & Clinical Medicine* 2021;8[4]: 39-52.
25. Jagadish U, Kumar K V, Shanthappa AH. Functional Outcome of Distal Humerus Fractures Treated with Open Reduction and Internal Fixation with Bicondylar Plating in a Tertiary Care Setting. *Cureus*. 2023 Jan 9;15[1]: e33540. doi: 10.7759/cureus.33540.
26. Patel SS, Mir HR, Horowitz E, Smith C, Ahmed AS, Downes K, Nydick JA. ORIF of Distal Humerus Fractures with Modern Pre-contoured Implants is Still Associated with a High Rate of Complications. *Indian J Orthop*. 2020 Apr 24;54[5]:570-579. doi: 10.1007/s43465-020-00124-4.

IJMA



INTERNATIONAL JOURNAL OF MEDICAL ARTS

Volume 7, Issue 10 (October 2025)



<http://ijma.journals.ekb.eg/>

P-ISSN: 2636-4174

E-ISSN: 2682-3780